

Algorithms for stochastic nonconvex and nonsmooth optimization

Courtney Paquette

University of Waterloo

Applied Math and Analysis Seminar, Duke University
January 25, 2019

Research Directions

(1). Thesis Work

- ▶ computational complexity
- ▶ nonsmooth analysis of eigenvalues
- ▶ composite nonlinear models ($h \circ c$)
- ▶ statistical guarantees for nonconvex problems

(2). Post doc

- ▶ stochastic optimization
- ▶ constrained conjugate gradient

Research Directions

(1). Thesis Work

- ▶ computational complexity
- ▶ nonsmooth analysis of eigenvalues
- ▶ composite nonlinear models ($h \circ c$)
- ▶ **statistical guarantees for nonconvex problems**

(2). Post doc

- ▶ **stochastic optimization**
- ▶ constrained conjugate gradient

- (1). Local search for non-smooth and non-convex problems
- (2). Adaptive line search for stochastic optimization

Local search for non-smooth and non-convex problems

Joint work with D. Davis (Cornell), D. Drusvyatskiy (U. Washington),
and K. MacPhee (U. Washington)

Why study nonsmooth and nonconvex optimization?

$$\min_x g(x)$$

Nonsmooth and nonconvex losses arise often...

- Structure (sparsity), robustness (outliers), stability (better conditioning)

Common problem class: $(\text{convex}) \circ (\text{smooth})$

(Fletcher '80, Powell '83, Burke '85, Wright '90, Lewis-Wright '08, Cartis-Gould-Toint '11)

Example: Robust Phase Retrieval

Problem: Find $x \in \mathbb{R}^d$ such that

$$(a_i^T x)^2 \approx b_i \quad a_1, \dots, a_m \in \mathbb{R}^d, \quad b_1, \dots, b_m \in \mathbb{R}.$$

Composite formulation:

$$\min_x g(x) := \frac{1}{m} \sum_{i=1}^m |(a_i^T x)^2 - b_i|$$

Assumptions:

$$m \gtrsim d, \quad b_i = (a_i^T \bar{x})^2 \text{ for some } \bar{x} \in \mathbb{R}^d, \quad a_i \sim N(0, I_d) \text{ and independent}$$

Example: Robust Phase Retrieval

Problem: Find $x \in \mathbb{R}^d$ such that

$$(a_i^T x)^2 \approx b_i \quad a_1, \dots, a_m \in \mathbb{R}^d, \quad b_1, \dots, b_m \in \mathbb{R}.$$

Composite formulation:

$$\min_x g(x) := \frac{1}{m} \sum_{i=1}^m |(a_i^T x)^2 - b_i|$$

Assumptions:

$$m \gtrsim d, \quad b_i = (a_i^T \bar{x})^2 \text{ for some } \bar{x} \in \mathbb{R}^d, \quad a_i \sim N(0, I_d) \text{ and independent}$$

Consequences: $\exists$ constants $\rho, \tau > 0$ such that with probability $1 - e^{-cm}$

- **Weakly-convex:** (Duchi-Ruan '17)

$$x \mapsto g(x) + \frac{\rho}{2} \|x\|_2^2 \quad \text{is convex}$$

- **Sharpness:** (Eldar-Mendelson '14)


$$g(x) \geq \tau \|\bar{x}\|_2 \operatorname{dist}(x, \{\pm \bar{x}\}).$$

Holds even when up to 1/2 the points are **corrupted!**

Intuition

$g(x) = \frac{1}{m} \sum_{i=1}^m |(a_i^T x)^2 - b_i|$ approximates the **population objective**:

$$g_P(x) = \mathbf{E}_{a \sim N}[|(a^T x)^2 - (a^T \bar{x})^2|]$$



Local search

$$\min_x g(x)$$

Strategy:

- Find a moderately accurate solution $\hat{x}$ at a low sample complexity cost
 - ▶ Available for: phase retrieval (Candès et al. '15), blind deconvolution (Ma et al. '17, Li et al. '18), matrix sensing (Boczar et al. '16)
- Refine $\hat{x}$ with a rapidly converging algorithm

Local search

$$\min_x g(x)$$

g is μ -sharp and ρ -weakly convex

Strategy:

- Find a moderately accurate solution $\hat{x}$ at a low sample complexity cost
 - ▶ Available for: phase retrieval (Candès et al. '15), blind deconvolution (Ma et al. '17, Li et al. '18), matrix sensing (Boczar et al. '16)
- Refine $\hat{x}$ with a rapidly converging algorithm

Is there a generic gradient-based **local search procedure** for nonsmooth and nonconvex problems?

Stationary points

- **Subdifferential** $\partial g(x)$ consists of $v \in \mathbb{R}^d$ such that

$$g(y) \geq g(x) + \langle v, y - x \rangle + o(\|y - x\|), \quad \text{as } y \rightarrow x$$

- Under weak-convexity,

$$x \text{ stationary} \Leftrightarrow \inf_{\|v\|=1} g'(x, v) \geq 0 \Leftrightarrow 0 \in \partial g(x).$$

Remark:

$$\partial(h \circ c)(x) = \nabla c(x)^T \partial h(c(x))$$

Stationary points

- **Subdifferential** $\partial g(x)$ consists of $v \in \mathbb{R}^d$ such that

$$g(y) \geq g(x) + \langle v, y - x \rangle + o(\|y - x\|), \quad \text{as } y \rightarrow x$$

- Under weak-convexity,

$$x \text{ stationary} \Leftrightarrow \inf_{\|v\|=1} g'(x, v) \geq 0 \Leftrightarrow 0 \in \partial g(x).$$


Remark:

$$\partial(h \circ c)(x) = \nabla c(x)^T \partial h(c(x))$$

Lemma (Davis-Drusvyatskiy-MacPhee-P)

No extraneous stationary points of g lie in the tube:

$$\mathcal{T} := \left\{ x \in \mathbb{R}^d : \text{dist}(x; S) < \frac{\mu}{\rho} \right\}$$



Meta-Theorem:

Simple algorithms for **sharp** and **weakly convex** functions converge rapidly

Meta-Theorem:

Simple algorithms for **sharp** and **weakly convex** functions converge rapidly

$$\min_x g(x) := h(c(x)), \quad (\text{convex}) \circ (\text{smooth})$$

Prox-linear method:

$$x^+ := \operatorname{argmin}_y \left\{ h(c(x) + \nabla c(x)(y - x)) + \frac{\rho}{2} \|y - x\|^2 \right\}$$

(Burke '85, '91, Fletcher '82, Powell '84, Wright '90, Yuan '83, Cartis-Gould-Toint '11)

Eg: proximal gradient, Levenberg-Marquardt

Subgradient method:

$$x^+ = x - \left(\frac{g(x) - \inf g}{\|v\|^2} \right) v \quad \text{where } v \in \partial g(x)$$

Prox-linear method

$$x^+ = \operatorname{argmin}_y \{h(c(x) + \nabla c(x)(y - x)) + \frac{\rho}{2} \|y - x\|^2\}$$

Thm: (Burke-Ferris '93)

Suppose that $g = h \circ c$ is

- ρ -weakly convex
- μ -sharp
- $\operatorname{dist}(x_0, S) \leq \frac{\mu}{2\rho}$

Then

$$\operatorname{dist}(x_{k+1}, S) \leq \frac{\mu}{\rho} \cdot \left(\frac{1}{2}\right)^{2^k} \quad \text{for all } k.$$

Global convergence guarantees (Drusvyatskiy-P, Math. Program '16)

Eg: phase retrieval

- $\frac{\mu}{\rho}$ is **dimension independent** w.h.p. (Eldar-Mendelson '14, Duchi-Ruan '17)

Subgradient Methods

Polyak subgradient method:

$$x^+ = x - \left(\frac{g(x) - \inf g}{\|v\|^2} \right) v \quad \text{where } v \in \partial g(x).$$

Thm: (Polyak '67, Davis-Drusvyatskiy-MacPhee-P '17)

Suppose that g is

- ρ -weakly convex
- L -Lipschitz
- μ -sharp
- $\text{dist}(x_0, S) \leq \frac{\mu}{2\rho}$

Then

$$\frac{\text{dist}(x_{k+1}, S)}{\text{dist}(x_k, S)} \leq \sqrt{1 - \left(\frac{\mu}{L\sqrt{2}} \right)^2}, \quad \text{for all } k.$$

Eg: phase retrieval

- $\frac{\mu}{\rho}, \frac{\mu}{L}$ are **dimension independent** w.h.p. (Eldar-Mendelson '14, Davis-Drusvyatskiy-MacPhee-P '17)




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 1.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 2.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 3.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 4.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 5.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 6.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 7.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 8.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 9.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 10.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 11.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 12.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 13.





Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 14.




Figure: $(d, m) \approx (2^{23}, 2^{24})$. Iteration 15.




Figure: Convergence plot (iterates vs. $\|x_k - \bar{x}\|/\|\bar{x}\|$).

Research Program:

Physical/Statistical well-posedness $\Rightarrow$ $(\text{convex}) \circ (\text{smooth})$ with “sharpness”-like conditions $\Rightarrow$ Simple algorithms converge rapidly

- Robust phase retrieval (Duchi-Ruan '17, Davis-Drusvyatskiy-MacPhee-P '17)
- Blind deconvolution/bi-convex sensing (Ling-Strohmer '15, Ahmed et al. '14)
- Covariance estimation (Chen et. al '15)
- Robust PCA (Candes et al. '11, Chandrasekaran et al. '11, Netrapalli et al. '14)
- Non-negative matrix factorization (Lee-Seung '99, Donoho-Stodden '03)

Adaptive line search method
for **smooth** stochastic optimization

Joint work with K. Scheinberg

Stochastic optimization

Central task in machine learning

$$\min_x f(x) = \mathbf{E}_{\xi \sim P}[\tilde{f}(x; \xi)]$$

Stochastic gradient descent (SGD):

$$\left\{ \begin{array}{l} \text{Pick } \xi \sim P \\ \text{Set } x_{k+1} = x_k - \alpha_k \nabla \tilde{f}(x_k; \xi) \end{array} \right\}$$

- **Major drawback:** stepsize, α_k , requires lots of tuning

Stochastic optimization

Central task in machine learning

$$\min_x f(x) = \mathbf{E}_{\xi \sim P}[\tilde{f}(x; \xi)]$$

Stochastic gradient descent (SGD):

$$\left\{ \begin{array}{l} \text{Pick } \xi \sim P \\ \text{Set } x_{k+1} = x_k - \alpha_k \nabla \tilde{f}(x_k; \xi) \end{array} \right\}$$

- **Major drawback:** stepsize, α_k , requires lots of tuning

Deterministic setting: Use **line search techniques**

Question:

Can the line search technique be adapted to the **stochastic** setting?

(Deterministic) Backtracking Line Search

Classical problem

$$\min_{x \in \Omega} f(x)$$

$f : \Omega \rightarrow \mathbf{R}$ with L -Lipschitz gradient

Gradient descent: $x_{k+1} = x_k - \alpha \nabla f(x_k), \quad \alpha \in (0, 1/L]$

(Deterministic) Backtracking Line Search

Classical problem

$$\min_{x \in \Omega} f(x)$$

$f : \Omega \rightarrow \mathbf{R}$ with L -Lipschitz gradient

Gradient descent: $x_{k+1} = x_k - \alpha \nabla f(x_k)$, $\alpha \in (0, 1/L]$

Backtracking Line Search Algorithm

- Compute $f(x_k)$ and $\nabla f(x_k)$
- Check sufficient decrease (Armijo '66)

$$f(x_k - \alpha_k \nabla f(x_k)) \leq f(x_k) - \theta \alpha_k \|\nabla f(x_k)\|^2$$

- Successful: $x_{k+1} = x_k - \alpha_k \nabla f(x_k)$ and increase $\alpha_k \Rightarrow \alpha_{k+1} = \gamma^{-1} \alpha_k$
- Unsuccessful: $x_{k+1} = x_k$ and decrease $\alpha_k \Rightarrow \alpha_{k+1} = \gamma \alpha_k$

(Deterministic) Backtracking Line Search

Classical problem

$$\min_{x \in \Omega} f(x)$$

$f : \Omega \rightarrow \mathbb{R}$ with L -Lipschitz gradient

Gradient descent: $x_{k+1} = x_k - \alpha \nabla f(x_k)$, $\alpha \in (0, 1/L]$

Backtracking Line Search Algorithm

- Compute $f(x_k)$ and $\nabla f(x_k)$
- Check sufficient decrease (Armijo '66)

$$\underbrace{f(x_k - \alpha_k \nabla f(x_k))}_{\text{function value at next step}} \leq \underbrace{f(x_k) - \theta \alpha_k \|\nabla f(x_k)\|^2}_{\text{linearization of } f \text{ at current step}}$$

- Successful: $x_{k+1} = x_k - \alpha_k \nabla f(x_k)$ and increase $\alpha_k \Rightarrow \alpha_{k+1} = \gamma^{-1} \alpha_k$
- Unsuccessful: $x_{k+1} = x_k$ and decrease $\alpha_k \Rightarrow \alpha_{k+1} = \gamma \alpha_k$

Question

Can the line search technique be adapted to **stochastic** setting using only **knowable** quantities?

Knowable quantities: e.g. bound on variance of $\nabla \tilde{f}, \tilde{f}$

Related works

Line search & heuristics Previous work requires: $\nabla f(x), \alpha_k \rightarrow 0$

- Bollapragada, Byrd, and Nocedal; “Adaptive sampling strategies for stochastic optimization” (to appear in SIOPT 2017)
- Friedlander and Schmidt; “Hybrid deterministic-stochastic methods for data fitting” (2012, SIAM Sci. Comput)
- Mahsereci and Hennig; “Probabilistic line search for stochastic optimization” (JMLR 2018; NIPS 2015)

Stochastic backtracking line search

- Compute **stochastic** estimates $\underbrace{g_k}_{\nabla f(x_k)}$, $\underbrace{f_k}_{f(x_k)}$, and $\underbrace{f_k^+}_{f(x_k - \alpha_k g_k)}$

- Check sufficient decrease (**Armijo '66**)

$$f_k^+ \leq f_k - \theta \alpha_k \|g_k\|^2$$

- Successful: $x_{k+1} = x_k - \alpha_k g_k$ and **increase** $\alpha_k \Rightarrow \alpha_{k+1} = \gamma^{-1} \alpha_k$
- Unsuccessful: $x_{k+1} = x_k$ and **decrease** $\alpha_k \Rightarrow \alpha_{k+1} = \gamma \alpha_k$

Stochastic backtracking line search

- Compute **stochastic** estimates $\underbrace{g_k}_{\nabla f(x_k)}$, $\underbrace{f_k}_{f(x_k)}$, and $\underbrace{f_k^+}_{f(x_k - \alpha_k g_k)}$
- Check sufficient decrease (**Armijo '66**)
$$f_k^+ \leq f_k - \theta \alpha_k \|g_k\|^2$$
- Successful: $x_{k+1} = x_k - \alpha_k g_k$ and **increase** $\alpha_k \Rightarrow \alpha_{k+1} = \gamma^{-1} \alpha_k$
- Unsuccessful: $x_{k+1} = x_k$ and **decrease** $\alpha_k \Rightarrow \alpha_{k+1} = \gamma \alpha_k$

Challenges

$$f_k^+ \leq f_k - \theta \alpha_k \|g_k\|^2 \quad \stackrel{??}{\Rightarrow} \quad f(x_k - \alpha_k g_k) \leq f(x_k) - \theta \alpha_k \|\nabla f(x_k)\|^2$$

- Bad function estimates may $\uparrow$ objective value

Increase at most $\alpha_k^2 \|g_k\|^2$

- Stepsizes, α_k , become arbitrarily small

Stochastic line search

Algorithm

- Compute **random** estimate of the gradient, g_k
- Compute **random** estimate of $f_k \approx f(x_k)$ and $f_k^+ \approx f(x_k - \alpha_k g_k)$
- Check the **stochastic** sufficient decrease

$$f_k^+ \leq f_k - \theta \alpha_k \|g_k\|^2$$

- Successful: $x_{k+1} = x_k - \alpha_k g_k$ and $\alpha_k \uparrow \Rightarrow \alpha_{k+1} = \gamma^{-1} \alpha_k$

- ▶ Reliable step: If $\alpha_k \|g_k\|^2 \geq \delta_k^2$, $\uparrow \delta_k \Rightarrow \delta_{k+1}^2 = \gamma^{-1} \delta_k^2$
 - ▶ Unreliable step: If $\alpha_k \|g_k\|^2 < \delta_k^2$, $\downarrow \delta_k \Rightarrow \delta_{k+1}^2 = \gamma \delta_k^2$

- Unsuccessful: $x_{k+1} = x_k$, **decrease** α_k , and **decrease** δ_k
 $\Rightarrow \alpha_{k+1} = \gamma \alpha_k$ and $\delta_{k+1}^2 = \gamma \delta_k^2$.

Randomness assumptions

- **Accurate gradient** g_k **w/ prob.** p_g :

$$\Pr(\|g_k - \nabla f(x_k)\| \leq \alpha_k \|g_k\|) \geq p_g$$

- **Accurate function estimates** f_k **and** f_k^+ **w/ prob.** p_f :

$$\Pr(|f(x_k) - f_k| \leq \alpha_k^2 \|g_k\|^2$$

$$\text{and} \quad |f(x_k - \alpha_k g_k) - f_k^+| \leq \alpha_k^2 \|g_k\|^2) \geq p_f$$

Randomness assumptions

- **Accurate gradient** g_k w/ prob. p_g :

$$\Pr(\|g_k - \nabla f(x_k)\| \leq \alpha_k \|g_k\|) \geq p_g$$

- **Accurate function estimates** f_k and f_k^+ w/ prob. p_f :

$$\Pr(|f(x_k) - f_k| \leq \alpha_k^2 \|g_k\|^2$$

$$\text{and } |f(x_k - \alpha_k g_k) - f_k^+| \leq \alpha_k^2 \|g_k\|^2) \geq p_f$$

- **Variance condition**

$$\mathbb{E}[|f_k - f(x_k)|^2] \leq \theta^2 \delta_k^4 \quad (\text{same for } f_k^+).$$

Question: How to choose these probabilities (p_f, p_g) large enough?

$p_f, p_g \geq 1/2$ at least, but p_f should be large.

Satisfying randomness assumptions

$$\min_x f(x) = \mathbf{E}_{\xi \sim P}[\tilde{f}(x; \xi)]$$

and bound on variance

$$\mathbf{E}_{\xi \sim P}(\|\nabla \tilde{f}(x, \xi) - \nabla f(x)\|^2) \leq V_g, \quad \mathbf{E}_{\xi \sim P}(|\tilde{f}(x; \xi) - f(x)|^2) \leq V_f.$$

Satisfying randomness assumptions

$$\min_x f(x) = \mathbf{E}_{\xi \sim P}[\tilde{f}(x; \xi)]$$

and bound on variance

$$\mathbf{E}_{\xi \sim P}(\|\nabla \tilde{f}(x, \xi) - \nabla f(x)\|^2) \leq V_g, \quad \mathbf{E}_{\xi \sim P}(|\tilde{f}(x; \xi) - f(x)|^2) \leq V_f.$$

Example: sampling

$$g_k = \frac{1}{|S_g|} \sum_{i \in S_g} \nabla \tilde{f}(x_k; \xi_i), \quad f_k = \frac{1}{|S_f|} \sum_{i \in S_f} \tilde{f}(x_k; \xi_i).$$

How many samples do we need?

Satisfying randomness assumptions

$$\min_x f(x) = \mathbf{E}_{\xi \sim P}[\tilde{f}(x; \xi)]$$

and bound on **variance**

$$\mathbf{E}_{\xi \sim P}(\|\nabla \tilde{f}(x, \xi) - \nabla f(x)\|^2) \leq V_g, \quad \mathbf{E}_{\xi \sim P}(|\tilde{f}(x; \xi) - f(x)|^2) \leq V_f.$$

Example: sampling

$$g_k = \frac{1}{|S_g|} \sum_{i \in S_g} \nabla \tilde{f}(x_k; \xi_i), \quad f_k = \frac{1}{|S_f|} \sum_{i \in S_f} \tilde{f}(x_k; \xi_i).$$

How many samples do we need?

Chebyshev Inequality

$$|S_g| \approx \tilde{O} \left(\frac{V_g}{\alpha_k^2 \|g_k\|^2} \right), \quad |S_f| \approx \tilde{O} \left(\max \left\{ \frac{V_f}{\alpha_k^4 \|g_k\|^4}, \frac{V_f}{\delta_k^4} \right\} \right)$$

Convergence result

Key observations

- $\Phi_k = \underbrace{\nu(f(x_k) - \inf f) + (1 - \nu)\alpha_k \|\nabla f(x_k)\|^2}_{\text{balance each other}} + (1 - \nu)\theta\delta_k^2$
- $T_\varepsilon = \inf\{k \geq 0 : \|\nabla f(x_k)\| < \varepsilon\}$

Convergence result

Key observations

- $\Phi_k = \underbrace{\nu(f(x_k) - \inf f) + (1 - \nu)\alpha_k \|\nabla f(x_k)\|^2}_{\text{balance each other}} + (1 - \nu)\theta\delta_k^2$
- $T_\varepsilon = \inf\{k \geq 0 : \|\nabla f(x_k)\| < \varepsilon\}$

Thm: (P-Scheinberg '18) If $p_g p_f > 1/2$ and p_f sufficiently large, then

$$\mathbf{E}[\Phi_{k+1} - \Phi_k | \text{past}] \leq - \left(\alpha_k \|\nabla f(x_k)\|^2 + \theta \delta_k^2 \right)$$

and consequently,

$$\mathbf{E}[T_\varepsilon] \leq \mathcal{O} \left(\frac{1}{\varepsilon^2} \right).$$

Convergence result

Key observations

- $\Phi_k = \underbrace{\nu(f(x_k) - \inf f) + (1 - \nu)\alpha_k \|\nabla f(x_k)\|^2}_{\text{balance each other}} + (1 - \nu)\theta\delta_k^2$
- $T_\varepsilon = \inf\{k \geq 0 : \|\nabla f(x_k)\| < \varepsilon\}$

Thm: (P-Scheinberg '18) If $p_g p_f > 1/2$ and p_f sufficiently large, then

$$\mathbf{E}[\Phi_{k+1} - \Phi_k | \text{past}] \leq - \left(\alpha_k \|\nabla f(x_k)\|^2 + \theta \delta_k^2 \right)$$

and consequently,

$$\mathbf{E}[T_\varepsilon] \leq \mathcal{O} \left(\frac{1}{\varepsilon^2} \right).$$

Proof sketch

- α_k behave like random walk with $p = p_g p_f$
- $\{\Phi_k\}$ supermartingales with stopping time T_ε
- Optional stopping time and Wald's identity

Convex case

Assumptions:

- f is **convex** and $\|\nabla f(x)\| \leq L_f$ for all $x \in \Omega$
- $\|x - x^*\| \leq D$ for all $x \in \Omega$

Key observation:

$$\Phi_k = \frac{1}{\nu\varepsilon} - \frac{1}{\Psi_k}$$

where $\Psi_k = \nu(f(x_k) - \inf f) + (1 - \nu)\alpha_k \|\nabla f(x_k)\|^2 + (1 - \nu)\theta\delta_k^2$

Stopping time: $T_\varepsilon = \inf\{k : f(x_k) - \inf f < \varepsilon\}$

Convergence rate, convex (P-Scheinberg '18)

If $p_g p_f > 1/2$ and p_f sufficiently large, then

$$\mathbf{E}[T_\varepsilon] \leq \mathcal{O}\left(\frac{1}{\varepsilon}\right)$$

μ -strongly convex case

Key observation:

$$\Phi_k = \log(\Psi_k) - \log(\nu\varepsilon)$$

where $\Psi_k = \nu(f(x_k) - \inf f) + (1 - \nu)\alpha_k \|\nabla f(x_k)\|^2 + (1 - \nu)\theta\delta_k^2$



Convergence rate, strongly convex (P-Scheinberg '18)

If $p_g p_f > 1/2$ and p_f sufficiently large, then

$$\mathbf{E}[T_\varepsilon] \leq \mathcal{O}\left(\frac{L}{\mu} \log\left(\frac{1}{\varepsilon}\right)\right)$$

Preliminary results

$$\min_{\theta} \frac{1}{m} \sum_{i=1}^m \log(1 + \exp(-y_i(\theta^T x_i))) + \frac{\lambda}{2} \|\theta\|^2$$



Open questions and extensions

Conclusions

- General framework for convergence results
- Convergence analysis (nonconvex, convex, and strongly convex) for a line search algorithm with gradient descent.

Open questions and extensions

Conclusions

- General framework for convergence results
- Convergence analysis (nonconvex, convex, and strongly convex) for a line search algorithm with gradient descent.

Applications of the stochastic process

- Line search, trust region methods (Blanchet, Cartis, Menickelly, Scheinberg '17), and cubic regularization?
- Extensions into 2nd order stochastic methods with Hessian guarantees?

Open problems

- Finding a good practical stochastic line search for machine learning; sampling procedure too conservative
- Extending line search procedure to stochastic Wolfe conditions (BFGS)

References

Davis, D., Drusvyatskiy, D., MacPhee, K., and Paquette, C. (2018).

Subgradient methods for sharp, weakly convex functions.

J. Optim. Theory App.

Davis, D., Drusvyatskiy, D., and Paquette, C. (2017).

The nonsmooth landscape of phase retrieval.

arXiv:1711.03247.

Drusvyatskiy, D. and Paquette, C. (2018).

Efficiency of minimizing compositions of convex functions and smooth maps.

Math. Program.

Paquette, C. and Scheinberg, K. (2017).

A Stochastic Line Search Method with Convergence Rate Analysis.

arXiv: 1807.07994.